

표 1 . 신경두경부 핵심질문3 근거표

핵심질문 3

문헌정보	연구유형	대상자 수	문헌 질 KCIG
ACR Appropriateness Criteria; hearing loss and/or vertigo			
ACR Appropriateness Criteria; hearing loss and/or vertigo			
Fortnum H et al: The role of magnetic resonance imaging in the identification of suspected acoustic neuroma: a systematic review of clinical and cost-effectiveness and natural history. Health Technology Assessment 13:1-154, 2009			1
Newton JR et al: Magnetic resonance imaging screening in acoustic neuroma, Am J Otolaryngol 31:217-220, 2010			3
Annesely-Williams DJ et al: Magnetic resonance imaging in the investigation of sensorineural hearing loss: is contrast enhancement still necessary? J Laryngol Otol 115:14-21 2001			3
Schmalbrock P et al: Assessment of internal auditory canal tumors: a comparison of contrast-enhanced T1-weighted and steady-state T2-weighted gradient echo MR imaging. AJNR 20:1207-1213, 1999			3
Naganawa S et al: MR imaging of the inner ear: comparison of a three-dimensional fast spin-echo sequence with use of a dedicated quadrature-surface coil with gadolinium-enhanced spoiled gradient-recalled sequence. Radiology 208:679-685, 1998			3
Hermans R et al: MRI screening for acoustic neuroma without gadolinium: value of 3DFT-CISS sequence. Neuroradiology 39:593-598, 1997			4